


## Israeli People Committee's Report Find Catastrophic Side Effects Of Pfizer Vaccine To Every System In Human Body

### Description

via [GreatGameIndia](#)

The Israeli People Committee (IPC), a civilian body made of leading Israeli health experts, has published its April report into the [Pfizer vaccine's side effects indicating damage to almost every system in the human body](#). If the findings by IPC are genuine, then Pfizer vaccine is linked to more deaths in Israel than AstraZeneca's in the whole of Europe. The findings are catastrophic on every possible level. This is a detailed report that highlights the most devastating findings.



The verdict of the [Israeli People Committee](#) is that “there has never been a vaccine that has harmed as many people.” The report is long and detailed (read full report below).

*“We received 288 death reports in proximity to vaccination (90% up to 10 days after the vaccination), 64% of those were men.”*

Yet the report states, “according to data provided by the Ministry of Health, only 45 deaths in Israel were vaccine related.”

If these are the genuine numbers, then Israel has failed to report on its experimental results genuinely.

We have been hearing a lot about the rare side effects of the AstraZeneca vaccine and more than 300 cases of blood clots found in Europe.

German scientists have found the exact [2 step process how the AstraZeneca COVID-19 vaccine causes blood clots](#) in recipients. They describe a series of events that has to happen in the body before the vaccines create these large clots.

If the findings by IPC are genuine, then Pfizer vaccine is linked to more deaths in Israel than AstraZeneca's in the whole of Europe.

Meanwhile, the US CDC and FDA have lifted their recommended pause on use of Johnson & Johnson's coronavirus vaccine with a condition that it will now include a safety label [warning that its vaccine comes with blood clot risks](#).

*"According to Central Bureau of Statistics data during January-February 2021, at the peak of the Israeli mass vaccination campaign, there was a 22% increase in overall mortality in Israel compared with the previous year."*

*"In fact, January-February 2021 have been the deadliest months in the last decade, with the highest overall mortality rates compared to corresponding months in the last 10 years."*

The IPC finds that "amongst the 20-29 age group the increase in overall mortality has been most dramatic. In this age group, we detect an increase of 32% in overall mortality in comparison with previous year."

*"Statistical analysis of information from the Central Bureau of Statistics, combined with information from the Ministry of Health, leads to the conclusion that the mortality rate amongst the vaccinated is estimated at about 1: 5000 (1: 13000 at ages 20-49, 1: 6000 at ages 50-69, 1: 1600 at ages 70+)."*

*"According to this estimate, it is possible to estimate the number of deaths in Israel in proximity of the vaccine, as of today, at about 1000-1100 people."*

If this is a genuine statistical analysis, then numbers reported by health authorities of Israel are misleading to a very great extent.

*"There is a high correlation between the number of people vaccinated per day and the number of deaths per day, in the range of up to 10 days, in all age groups."*

*"Ages 20-49 – a range of 9 days from the date of vaccination to mortality, ages 50-69 – 5 days from the date of vaccination to mortality, ages 70 and up – 3 days from the date of*

*vaccination to mortality.”*

The IPC also reveals that the “the risk of mortality after the second vaccine is higher than the risk of mortality after the first vaccine.”

This is not only about the death risk, as per the reports of IPC, “as of the date of publication of the report, 2066 reports of side effects have accumulated in the Civil Investigation Committee and the data continue to come in.

*These reports indicate damage to almost every system in the human body. Our analysis found a relatively high rate of heart-related injuries.*

*26% of all cardiac events occurred in young people up to the age of 40, with the most common diagnosis in these cases being Myositis or Pericarditis.*

*Also, a high rate of massive vaginal bleeding, neurological damage, and damage to the skeletal and skin systems has been observed.*

*It should be noted that a significant number of reports of side effects are related, directly or indirectly, to Hypercoagulability (infarction), Myocardial infarction, stroke, miscarriages, impaired blood flow to the limbs, pulmonary embolism.”*

*Read the summary of the Israeli People Committee report in English below.*

--  
Page 1 / 6

--  
Zoom 100%

*Read the full Israeli People Committee report in Hebrew below.*